Kyneton to Heathcote regions

Common Eastern Froglet (Crinia signifera)

Breeding: All year

Description: Common, widespread.

Most likely species heard.

Up to 3cm long.

Variable colouration and patterning, from brown to red to mustard, with or without stripes or spots.





Eastern Banjo Frog, Pobblebonk (Limnodynastes dumerilii)

Breeding: Spring to Autumn

Description: Distinctive call.

Large frog, up to 7.5cm long.

Brown or grey-ish back, often with orange or yellow mottling on sides.

Pale stripe under the eye to the shoulder.







Kyneton to Heathcote regions

Eastern Sign-Bearing Froglet (Crinia parinsignifera)

Breeding: All year (except mid-Winter)

Description: Small, up to 2.5cm long.

Back can be grey, brown or sandy-gold, with or without stripes.

Often a dark patch between the eyes and dark bars on the legs.

Cream-coloured belly with several small light brown spots.





Spotted Marsh Frog (Limnodynastes tasmaniensis)

Breeding: Spring to Autumn

Description: Up to ~5cm long.

Grey-brown or olive-green back with darker 'spots'.

Often a pale cream or red stripe down the middle of the back.

Cream-coloured stripe from under the eye to the top of the arm





Kyneton to Heathcote regions

Peron's / Laughing Tree Frog (Litoria peronii)

Breeding: Spring to Autumn

Description: Up to 7cm long.

Grey-to-light-brown back, with or without darker mottling and tiny emerald-green spots.

Thin black line from behind the eye to the top of the arm.

Bright yellow thighs with black patches.





Southern Brown Tree Frog (Litoria ewingii)

Breeding: All year (after rain)

Description: Up to 4.5cm long.

Cream, brown, or yellow-brown back, with a darker wide stripe along the middle.

Dark stripe from snout to past the arm. Often a white stripe from below the eye to the end of the upper lip.

Backs of thighs are reddish-orange.





Kyneton to Heathcote regions

Striped Marsh Frog (Limnodynastes peronii)

Breeding: Spring to Autumn

Description: Up to 7.5cm long.

Brown back with dark stripes, sometimes a cream or reddish stripe down the middle.

Pale stripe from under the eye to top of the arm.

Males have a pale yellow throat with brown mottling.



Bibron's Toadlet (Pseudophryne bibronii)

Breeding: All year (after rain)

Description: Up to 3cm long.

Dark grey, blue-grey or brown back, often with small black spots and a red-orange stripe down the lower half.

Sometimes a light brown or pale orange patch on the head.

Tops of shoulders and the back of thighs are yellow to orange.





Images sourced from frogid.net.au Information from frogid.net.au, frogs.org.au and ala.org.au.

Kyneton to Heathcote regions

Growling Grass Frog (Litoria raniformis)

Breeding: Spring to Autumn

Description: Up to 10cm long.

Green-brown back, with brown-bronze patches, and sometimes a pale stripe down the middle.

Cream-coloured stripe from behind the eye along the sides, and often a dark brown stripe from the nostril to the eye.

Groin and backs of the thighs are bright blue, sometimes with small, bright yellow patches.



Eastern Smooth Frog/Victorian Smooth Froglet

(Geocrinia victoriana)

Breeding: Summer to Autumn

Description: Up to 3.5cm long.

Brown-grey or red-brown back, often dark, yellow, white, or orange patches.

Often a dark horizontal stripe between the eyes, a small dark stripe behind the eye, and dark patches along the upper lip.

Groin and back of the lower legs sometimes have a pink or orange-brown patch. Legs sometimes have dark horizontal bars.





Images sourced from frogid.net.au Information from frogid.net.au, frogs.org.au and ala.org.au.